

THE GRILLAKIS-MACHEDON-MARGETIS SECOND ORDER CORRECTIONS TO MEAN FIELD EVOLUTION FOR WEAKLY INTERACTING BOSONS IN THE CASE OF 3-BODY INTERACTIONS

XUWEN CHEN

ABSTRACT. This note shows that, with a little modification, the results in [1] hold in the 3-body interactions case.

1. INTRODUCTION

Recently, a second-order correction to the usual tensor product (mean-field) approximation for the Hamiltonian evolution of a many-particle system with 2-body potential was found in Grillakis-Machedon-Margetis[1]. Here is their main result,

Theorem 1. [1] *Let ϕ be a smooth solution of the Hartree equation*

$$(1.1) \quad i\frac{\partial\phi}{\partial t} + \Delta\phi + (v' * |\phi|^2)\phi = 0$$

with initial conditions ϕ_0 and potential v' being even. Assuming the following:

- (1) *Let $k(t, x, y) \in L_s^2(dxdy)$ for a.e. t , solve the Grillakis-Machedon-Margetis equation*

$$(1.2) \quad (iu_t + ug^T + gu - (I + p)m) = (ip_t + [g, p] + u\overline{m})(1 + p)^{-1}u ,$$

where products in equation 1.2 mean compositions of operators, and

$$u(t, x, y) := \sinh(k) := k + \frac{1}{3!}k\bar{k}k + \dots ,$$

$$\delta(x - y) + p(t, x, y) := \cosh(k) := \delta(x - y) + \frac{1}{2!}k\bar{k} + \dots ,$$

$$g(t, x, y) := -\Delta_x\delta(x - y) - v(x - y)\phi(t, x)\overline{\phi}(t, y) - (v * |\phi|^2)(t, x)\delta(x - y) ,$$

$$m(t, x, y) := v(x - y)\overline{\phi}(t, x)\overline{\phi}(t, y) .$$

- (2) *Functions $f(t) := \|e^B[A, V']e^{-B}\Omega\|_{\mathcal{F}}$ and $g(t) := \|e^BV'e^{-B}\Omega\|_{\mathcal{F}}$ are locally integrable, where*

$$(1.3) \quad B(t) = \frac{1}{2} \int (k(t, x, y)a_x a_y - \bar{k}(t, x, y)a_x^* a_y^*) dx dy$$

and $\mathcal{F}$ the Fock space defined in [1].

The author thanks Matei Machedon for discussions related to this work.

(3) $\int d(t, x, x) dx$ is locally integrable in time, where

$$(1.4) \quad \begin{aligned} d(t, x, y) = & (i \sinh(k)_t + \sinh(k)g^T + g \sinh(k)) \overline{\sinh(k)} \\ & - (i \cosh(k)_t + [g, \cosh(k)]) \cosh(k) \\ & - \sinh(k) \overline{m} \cosh(k) - \cosh(k) m \overline{\sinh(k)} . \end{aligned}$$

Then, there exist real functions χ_0, χ_1 such that

$$(1.5) \quad \begin{aligned} & \|e^{-\sqrt{N}A(t)}e^{-B(t)}e^{-i\int_0^t(N\chi_0(s)+\chi_1(s))ds}\Omega - e^{itH'_N}e^{-\sqrt{N}A(0)}\Omega\|_{\mathcal{F}} \\ & \leq \frac{\int_0^t f(s)ds}{\sqrt{N}} + \frac{\int_0^t g(s)ds}{N} , \end{aligned}$$

where $\Omega = (1, 0, 0, \dots) \in \mathcal{F}$ is the vacuum state,

$$(1.6) \quad A(\phi) = a(\bar{\phi}) - a^*(\phi)$$

with a and a^* are the annihilation operator and creation operator, and the Hamiltonian

$$\begin{aligned} H'_N &= \int a_x^* \Delta a_x dx + \frac{1}{2N} \int v'(x-y) a_x^* a_y^* a_x a_y dx dy \\ &=: H_0 + \frac{1}{2N} V' . \end{aligned}$$

To prove Theorem 1, the key is to notice that, for ϕ and k satisfying the assumptions, there is some "nice" Hermitian $\tilde{L}$ such that

$$\frac{1}{i} \frac{\partial}{\partial t} \Psi = (\tilde{L} - N\chi_0 - \chi_1) \Psi ,$$

and

$$\|\tilde{L}\Omega\|_{\mathcal{F}} \leq N^{-1/2} \|e^B[A, V]e^{-B}\Omega\|_{\mathcal{F}} + N^{-1} \|e^B V e^{-B}\Omega\|_{\mathcal{F}} ,$$

where

$$\Psi(t) = e^{B(t)} e^{\sqrt{N}A(t)} e^{itH} e^{-\sqrt{N}A(0)} \Omega .$$

Then applying the energy estimate to

$$\left(\frac{1}{i} \frac{\partial}{\partial t} - \tilde{L} \right) (e^{i\int_0^t(N\chi_0(s)+\chi_1(s))ds} \Psi - \Omega) = \tilde{L}\Omega ,$$

would give the estimate 1.5.

With the same set up, this result can be extended to the case of 3-body interaction potentials.

2. MAIN RESULT

We consider the Fock space Hamiltonian $H_N : \mathcal{F} \rightarrow \mathcal{F}$ with the 3-body interaction potential defined by

$$(2.1) \quad \begin{aligned} H_N &= \int a_x^* \Delta a_x dx + \frac{1}{6N^2} \int v(x-y, y-z) a_x^* a_y^* a_z^* a_x a_y a_z dx dy \\ &=: H_0 + \frac{1}{6N^2} V , \end{aligned}$$

in correspondence to the Hamiltonian

$$H_{N,N} = \sum_{j=1}^n \Delta_{x_j} + \frac{1}{N^2} \sum_{i < j < k}^n v(x_i - x_j, x_j - x_k) .$$

We assume that v is symmetric in x , y , and z . As shown in (1.2) and (1.3) in Chen-Pavlović[2], every translation invariant 3-body potential can be written in the form $v(x - y, y - z)$. However, I don't assume $v \geq 0$ as in [2].

Theorem 2. *With the same assumptions as in theorem 1 with H'_N replaced by H_N , V' by V in formula 2.1. We change equation 1.1 to*

$$(2.2) \quad i \frac{\partial}{\partial t} \phi + \Delta \phi + \frac{1}{2} \phi \int v(x - y, y - z) |\phi(y)|^2 |\phi(z)|^2 dydz = 0$$

and

$$g(t, x, y) := -\Delta \delta_{12} - \left(\int v_{12,23} |\phi_3|^2 dz \right) \bar{\phi}_1 \phi_2 - \frac{1}{2} \left(\int v_{12,23} |\phi_2|^2 |\phi_3|^2 dydz \right) \delta_{12}$$

$$m(t, x, y) := \left(\int v_{12,23} |\phi_3|^2 dz \right) \bar{\phi}_1 \bar{\phi}_2$$

where $v_{12,23} |\phi_3|^2$ is an abbreviation for the product $v(x - y, y - z) |\phi(z)|^2$,

Further, we assume that the functions

$$h(t) := \|e^B[A, [A, V]]e^{-B}\Omega\|_{\mathcal{F}}$$

and

$$i(t) := \|e^B[A, [A, [A, V]]]e^{-B}\Omega\|_{\mathcal{F}}$$

are also locally integrable.

Then, there exist real functions χ_0, χ_1 such that

$$\begin{aligned} & \|e^{-\sqrt{N}A(t)}e^{-B(t)}e^{-i \int_0^t (N\chi_0(s) + \chi_1(s))ds}\Omega - e^{itH_N}e^{-\sqrt{N}A(0)}\Omega\|_{\mathcal{F}} \\ & \leq \frac{\int_0^t f(s)ds}{6N^{\frac{3}{2}}} + \frac{\int_0^t g(s)ds}{6N^2} + \frac{\int_0^t h(s)ds}{12N} + \frac{\int_0^t i(s)ds}{36N^{\frac{1}{2}}}. \end{aligned}$$

Proof. Similar to the proof of Theorem 1. □

Remark 1. Recall that, the equation for k is 1.2

$$(2.3) \quad (iu_t + ug^T + gu - (I + p)m) = (ip_t + [g, p] + u\bar{m})(1 + p)^{-1}u.$$

A derivation of this equation will be given in Section 4.

Remark 2. In Section 4, we will show that the $\tilde{L}$ in this case is

$$(2.4) \quad \begin{aligned} \tilde{L} = & H_0 + \int v_{12,23} |\phi_3|^2 \bar{\phi}_2 \phi_1 a_x^* a_y dx dy dz + \\ & \frac{1}{2} \int v_{12,23} |\phi_2|^2 |\phi_3|^2 a_x^* a_x dx dy dz \\ & - \int d(t, x, y) a_y^* a_x dx + \frac{1}{6N^2} e^B V e^{-B} + \\ & \frac{1}{6N^{3/2}} e^B [A, V] e^{-B} + \frac{1}{12N} e^B [A, [A, V]] e^{-B} \\ & + \frac{1}{36N^{\frac{1}{2}}} e^B [A, [A, [A, V]]] e^{-B}, \end{aligned}$$

where $d(t, x, y)$ is the same as equation 1.4.

3. THE NEW HARTREE EQUATION

In this section we derive the Hartree equation 2.2 for the one-particle wave function ϕ as needed in Theorem 2.

Lemma 1. *The following relations hold, where A denotes $A(\phi)$, and A, V are defined by formulas 1.6 and 2.1):*

$$\begin{aligned}
& [A, V] \\
&= 3 \int v(x-y, y-z) (\bar{\phi}(x) a_y^* a_z^* a_x a_y a_z + \phi(x) a_x^* a_y^* a_z^* a_y a_z) dx dy dz \\
& \quad [A, [A, V]] \\
&= 6 \int v(x-y, y-z) (\bar{\phi}(x) \bar{\phi}(y) a_z^* a_x a_y a_z + 2\phi(x) \bar{\phi}(y) a_x^* a_z^* a_y a_z \\
& \quad + \phi(x) \phi(y) a_x^* a_y^* a_z^* a_y a_z) dx dy dz \\
& \quad + 6 \int v(x-y, y-z) |\phi(x)|^2 a_y^* a_z^* a_y a_z dx dy dz \\
& \quad [A, [A, [A, V]]] \\
&= 36 \int v(x-y, y-z) |\phi(x)|^2 (\bar{\phi}(y) a_z^* a_y a_z + \phi(y) a_y^* a_z^* a_z) dx dy dz \\
& \quad + 6 \int v(x-y, y-z) (\bar{\phi}(x) \bar{\phi}(y) \bar{\phi}(z) a_x a_y a_z + \phi(x) \phi(y) \phi(z) a_x^* a_y^* a_z^*) dx dy dz \\
& \quad + 18 \int v(x-y, y-z) (\bar{\phi}(x) \bar{\phi}(y) \phi(z) a_z^* a_x a_y + \phi(x) \phi(y) \bar{\phi}(z) a_x^* a_y^* a_z) dx dy dz \\
& \quad [A, [A, [A, [A, V]]]] \\
&= 72 \int v(x-y, y-z) |\phi(x)|^2 (\bar{\phi}(y) \bar{\phi}(z) a_y a_z + \phi(y) \phi(z) a_y^* a_z^*) dx dy dz \\
& \quad + 144 \int v(x-y, y-z) |\phi(x)|^2 \bar{\phi}(y) \phi(z) a_z^* a_y dx dy dz \\
& \quad + 72 \int v(x-y, y-z) |\phi(x)|^2 |\phi(y)|^2 a_z^* a_z dx dy dz \\
& \quad [A, [A, [A, [A, [A, V]]]]] \\
&= 360 \int v(x-y, y-z) |\phi(x)|^2 |\phi(y)|^2 (\bar{\phi}(z) a_z + \phi(z) a_z^*) dx dy dz \\
& \quad [A, [A, [A, [A, [A, [A, V]]]]]] \\
&= 720 \int v(x-y, y-z) |\phi(x)|^2 |\phi(y)|^2 |\phi(z)|^2 dx dy dz
\end{aligned}$$

Proof. This is a direct calculation. $\square$

Now, we write $\Psi_1(t) = e^{\sqrt{N}A(t)} e^{itH} e^{-\sqrt{N}A(0)} \Omega$ for which we carry out the calculation in the spirit of equation (3.7) in Rodnianski-Schlein [3].

Proposition 1. *Say ϕ satisfies the Hartree equation*

$$i \frac{\partial}{\partial t} \phi + \Delta \phi + \frac{1}{2} \phi \int v(x-y, y-z) |\phi(y)|^2 |\phi(z)|^2 dy dz = 0$$

then $\Psi_1(t)$ satisfies

$$\begin{aligned} \frac{1}{i} \frac{\partial}{\partial t} \Psi_1(t) = & \left(H_N + \frac{1}{4!} \frac{1}{6} [A, [A, [A, [A, V]]]] + \frac{1}{6} N^{-3/2} [A, V] \right. \\ & + \frac{1}{12} N^{-1} [A, [A, V]] + \frac{1}{36} N^{-\frac{1}{2}} [A, [A, [A, V]]] \\ & \left. - \frac{N}{3} \int v(x-y, y-z) |\phi(x)|^2 |\phi(y)|^2 |\phi(z)|^2 dx dy dz \right) \Psi_1(t) . \end{aligned}$$

Proof. Applying the formulas

$$\left(\frac{\partial}{\partial t} e^{C(t)} \right) \left(e^{-C(t)} \right) = \dot{C} + \frac{1}{2!} [C, \dot{C}] + \frac{1}{3!} [C, [C, \dot{C}]] + \dots$$

$$e^C H e^{-C} = H + [C, H] + \frac{1}{2!} [C, [C, H]] + \dots$$

to $C = \sqrt{N}A$ we have

$$(3.1) \quad \frac{1}{i} \frac{\partial}{\partial t} \psi_1(t) = L_1 \psi_1 ,$$

where

$$\begin{aligned} L_1 = & \frac{1}{i} \left(\frac{\partial}{\partial t} e^{\sqrt{N}A(t)} \right) e^{-\sqrt{N}A(t)} + e^{\sqrt{N}A(t)} H_N e^{-\sqrt{N}A(t)} \\ = & \frac{1}{i} \left(N^{1/2} \dot{A} + \frac{N}{2} [A, \dot{A}] \right) + H_N + N^{1/2} [A, H_0] + \frac{N}{2!} [A, [A, H_0]] + \frac{1}{6} \left(N^{-3/2} [A, V] \right. \\ & + \frac{N^{-1}}{2!} [A, [A, V]] + \frac{N^{-\frac{1}{2}}}{3!} [A, [A, [A, V]]] \\ & + \frac{1}{4!} [A, [A, [A, [A, V]]]] + \frac{N^{\frac{1}{2}}}{5!} [A, [A, [A, [A, [A, V]]]] \\ & \left. + \frac{N}{6!} [A, [A, [A, [A, [A, [A, V]]]]] \right) . \end{aligned}$$

The Hartree equation 2.2 is equivalent to setting

$$(3.2) \quad \frac{1}{i} \dot{A} + [A, H_0] + \frac{1}{6} \frac{1}{5!} [A, [A, [A, [A, [A, V]]]] = 0 .$$

Or more explicitly, equation 3.2 is

$$a(i\overline{\phi}_t + \overline{\Delta\phi} + \frac{1}{2}\overline{\phi} \int v_{12,23} |\phi_2|^2 |\phi_3|^2 dy dz) + a^*(i\phi_t + \Delta\phi + \frac{1}{2}\phi \int v_{12,23} |\phi_2|^2 |\phi_3|^2 dy dz) = 0 ,$$

because of Lemma 1 and the fact that $[\Delta_x a_x, a_y^*] = (\Delta\delta)(x-y)$.

Thus

$$\frac{1}{i} [A, \dot{A}] + [A, [A, H_0]] + \frac{1}{5!} \frac{1}{6} [A, [A, [A, [A, [A, [A, V]]]]] = 0 ,$$

i.e 3.1 simplifies to

$$(3.3) \quad \begin{aligned} \frac{1}{i} \frac{\partial}{\partial t} \Psi_1(t) = & \left(H_N + \frac{1}{4!} \frac{1}{6} [A, [A, [A, [A, V]]]] + \frac{1}{6} N^{-3/2} [A, V] \right. \\ & + \frac{1}{12} N^{-1} [A, [A, V]] + \frac{1}{36} N^{-\frac{1}{2}} [A, [A, [A, V]]] \\ & \left. - \frac{N}{3} \int v(x-y, y-z) |\phi(x)|^2 |\phi(y)|^2 |\phi(z)|^2 dx dy dz \right) \Psi_1(t) . \end{aligned}$$

□

Write the last term as

$$-\frac{N}{3} \int v(x-y, y-z) |\phi(x)|^2 |\phi(y)|^2 |\phi(z)|^2 dx dy dz := -N\chi_0 ,$$

then the first two terms on the right-hand side of 3.3 are the main ones we need to consider, since the next four terms are $O\left(\frac{1}{N^2}\right)$, $O\left(\frac{1}{N^{\frac{3}{2}}}\right)$, $O\left(\frac{1}{N}\right)$, and $O\left(\frac{1}{N^{\frac{1}{2}}}\right)$.

In order to kill the terms involving "only creation operators" i.e $a_x^* a_y^*$ in $\frac{1}{4!} \frac{1}{6} [A, [A, [A, [A, V]]]]$, we introduce B (see 1.3) and let

$$\Psi = e_1^B \Psi_1$$

as in [1]. So we have

$$\frac{1}{i} \frac{\partial}{\partial t} \Psi = L\Psi ,$$

where

$$\begin{aligned} L = & \frac{1}{i} \left(\frac{\partial}{\partial t} e^B \right) e^{-B} + e^B L_1 e^{-B} \\ = & L_Q + \frac{1}{6N^2} e^B V e^{-B} + \frac{1}{6} N^{-3/2} e^B [A, V] e^{-B} + \frac{1}{12} N^{-1} e^B [A, [A, V]] e^{-B} \\ & + \frac{1}{36} N^{-\frac{1}{2}} e^B [A, [A, [A, V]]] e^{-B} - N\chi_0 , \end{aligned}$$

and

$$(3.4) \quad L_Q = \frac{1}{i} \left(\frac{\partial}{\partial t} e^B \right) e^{-B} + e^B \left(H_0 + \frac{1}{4!} \frac{1}{6} [A, [A, [A, [A, V]]]] \right) e^{-B} .$$

4. EQUATION FOR k

In this section, we derive the equation for k .

Let $\mathcal{I} : sp \rightarrow Quad$ be the Lie algebra isomorphism defined by $Q(d, k, l) = \mathcal{I}(S(d, k, l))$ in theorem 3 in [1], where sp is the infinite dimensional Lie algebra containing

$$S(d, k, l) = \begin{pmatrix} d & k \\ l & -d^T \end{pmatrix}$$

and k and l are symmetric, and $Quad$ is the Lie algebra consisting of quadratics of the form

$$\begin{aligned} Q(d, k, l) &:= \frac{1}{2} \begin{pmatrix} a_x & a_x^* \end{pmatrix} \begin{pmatrix} d & k \\ l & -d^T \end{pmatrix} \begin{pmatrix} -a_y^* \\ a_y \end{pmatrix} \\ &= - \int d(x, y) \frac{a_x a_y^* + a_y^* a_x}{2} dx dy + \frac{1}{2} \int k(x, y) a_x a_y dx dy \\ &\quad - \frac{1}{2} \int l(x, y) a_x^* a_y^* dx dy . \end{aligned}$$

Recall that

$$\begin{aligned} &\frac{1}{4!6} [A, [A, [A, [A, V]]]] \\ &= \frac{1}{2} \int v(x-y, y-z) |\phi(x)|^2 (\bar{\phi}(y) \bar{\phi}(z) a_y a_z + \phi(y) \phi(z) a_y^* a_z^*) dx dy dz \\ &\quad + \int v(x-y, y-z) |\phi(x)|^2 \bar{\phi}(y) \phi(z) a_z^* a_y dx dy dz \\ &\quad + \frac{1}{2} \int v(x-y, y-z) |\phi(x)|^2 |\phi(y)|^2 a_z^* a_z dx dy dz , \end{aligned}$$

we write

$$G = \begin{pmatrix} g & 0 \\ 0 & -g^T \end{pmatrix} \quad \text{and} \quad M = \begin{pmatrix} 0 & m \\ -\bar{m} & 0 \end{pmatrix}$$

with

$$g = -\Delta \delta_{12} - \left(\int v_{12,23} |\phi_3|^2 dz \right) \bar{\phi}_1 \phi_2 - \frac{1}{2} \left(\int v_{12,23} |\phi_2|^2 |\phi_3|^2 dy dz \right) \delta_{12}$$

and

$$m = \left(\int v_{12,23} |\phi_3|^2 dz \right) \bar{\phi}_1 \bar{\phi}_2 .$$

Though it's true that

$$B = \mathcal{I}(K) ,$$

for

$$(4.1) \quad K = \begin{pmatrix} 0 & k(t, x, y) \\ \bar{k}(t, x, y) & 0 \end{pmatrix} ,$$

it's not quite true that

$$H_0 + \frac{1}{4!} \frac{1}{6} [A, [A, [A, [A, V]]]] = \mathcal{I}(G + M) .$$

However, the commutators of $\mathcal{I}(G + M)$ and $H_0 + \frac{1}{4!} \frac{1}{6} [A, [A, [A, [A, V]]]]$ with B are the same as in the discussion in page 15 in [1].

Now, L_Q reads

$$\begin{aligned} L_Q &= \frac{1}{i} \left(\frac{\partial}{\partial t} e^B \right) e^{-B} + e^B \left(H_0 + \frac{1}{4!} \frac{1}{6} [A, [A, [A, [A, V]]]] \right) e^{-B} \\ &= H_G + \mathcal{I} \left(\left(\frac{1}{i} \frac{\partial}{\partial t} e^K \right) e^{-K} + [e^K, G] e^{-K} + e^K M e^{-K} \right) \\ &= H_G + \mathcal{I}(\mathcal{M}_1 + \mathcal{M}_2 + \mathcal{M}_3) , \end{aligned}$$

where

$$H_G = H_0 + \int v_{12,23} |\phi_3|^2 \bar{\phi}_2 \phi_1 a_x^* a_y dx dy dz + \frac{1}{2} \int v_{12,23} |\phi_2|^2 |\phi_3|^2 a_x^* a_x dx dy dz,$$

in this setting.

Then by the definition of the isomorphism $\mathcal{I}$, the coefficient of $a_x a_y$ is $-(\mathcal{M}_{1,12} + \mathcal{M}_{2,12} + \mathcal{M}_{3,12})$, and the coefficient of $a_x^* a_y^*$ is $(\mathcal{M}_{1,21} + \mathcal{M}_{2,21} + \mathcal{M}_{3,21})$. To write it explicitly:

$$\begin{aligned} (4.2) \quad & -(\mathcal{M}_{1,12} + \mathcal{M}_{2,12} + \mathcal{M}_{3,12}) \\ & = \overline{(\mathcal{M}_{1,21} + \mathcal{M}_{2,21} + \mathcal{M}_{3,21})} \\ & = (i \sinh(k)_t + \sinh(k) g^T + g \sinh(k)) \overline{\cosh(k)} - \\ & \quad (i \cosh(k)_t - [\cosh(k), g]) \sinh(k) \\ & \quad - \sinh(k) \overline{m} \sinh(k) - \cosh(k) m \overline{\cosh(k)}. \end{aligned}$$

Setting formula 4.2 to 0 gives the equation 2.3. Hence we have the following theorem similar to Corollary 1 in [1].

Theorem 3. *If ϕ and k solve 2.2 and 2.3, then the coefficients of $a_x a_y$ and $a_x^* a_y^*$ drop out and L_Q becomes*

$$\begin{aligned} L_Q = & H_0 + \int v_{12,23} |\phi_3|^2 \bar{\phi}_2 \phi_1 a_x^* a_y dx dy dz + \frac{1}{2} \int v_{12,23} |\phi_2|^2 |\phi_3|^2 a_x^* a_x dx dy dz \\ & - \int d(t, x, y) \frac{a_x a_y^* + a_y^* a_x}{2} dx dy, \end{aligned}$$

where d is given by formula 1.4 and the full operator reads

$$\begin{aligned} L = & H_0 + \int v_{12,23} |\phi_3|^2 \bar{\phi}_2 \phi_1 a_x^* a_y dx dy dz + \frac{1}{2} \int v_{12,23} |\phi_2|^2 |\phi_3|^2 a_x^* a_x dx dy dz - \\ & \int d(t, x, y) a_y^* a_x dx + \frac{1}{6N^2} e^B V e^{-B} + \frac{1}{6N^{3/2}} e^B [A, V] e^{-B} + \frac{1}{12N} e^B [A, [A, V]] e^{-B} \\ & + \frac{1}{36N^{\frac{1}{2}}} e^B [A, [A, [A, V]]] e^{-B} - N \chi_0 - \chi_1 \\ & := \tilde{L} - N \chi_0 - \chi_1, \end{aligned}$$

and

$$\begin{aligned} \chi_0 = & \frac{1}{3} \int v(x-y, y-z) |\phi(x)|^2 |\phi(y)|^2 |\phi(z)|^2 dx dy dz, \\ \chi_1(t) = & -\frac{1}{2} \int d(t, x, x) dx. \end{aligned}$$

This matches equation 2.4.

Notice that

$$\begin{aligned} \tilde{L} \Omega = & \left(\frac{1}{6N^2} e^B V e^{-B} + \frac{1}{6} N^{-3/2} e^B [A, V] e^{-B} + \right. \\ & \left. \frac{1}{12} N^{-1} e^B [A, [A, V]] e^{-B} + \frac{1}{36} N^{-\frac{1}{2}} e^B [A, [A, [A, V]]] e^{-B} \right) \Omega, \end{aligned}$$

in our case. This ends the proof of theorem 2.

The verification that the hypotheses of the main theorem are satisfied by some singular potentials will be provided in a subsequent article.

REFERENCES

- [1] Grillakis, M. G., Machedon, M, Margetis, D.: Second order corrections to mean field evolution for weakly interacting Bosons. I. Preprint math-ph arXiv:0904.0158
- [2] Chen, T., Pavlović, N.: The quintic NLS as the mean field limit of a Boson gas with three-body interactions. Preprint math-ph arXiv:0812.2740
- [3] Rodnianski, I., Schlein, B.: Quantum fluctuations and rate of convergence towards mean field dynamics. Preprint math-ph arXiv:0711.3087

UNIVERSITY OF MARYLAND, COLLEGE PARK
E-mail address: `chenxuwen@math.umd.edu`